

Evaluation Of The Sensory Qualities Of Ready To Serve Blends Of Aloe Vera (*Aloe barbadensis* Mill.)

Mirza Arslan Abid and Ms. Najma Yousaf Zahid

Department of Horticulture, Pir Mehr Ali Shah Arid Agriculture University Rawalpindi Islamabad, Punjab, Pakistan.

Abstract: Aloe vera has been used for its medicinal worth for several thousand years. Its use is also mentioned in the Bible where it was taken as the plant of immortality. The research was conducted at Department of Horticulture, Pir Mehr Ali Shah Arid Agriculture University Rawalpindi, Punjab, Pakistan in year 2013-14. The main objectives of this study are to develop the economical Aloe drink using available resources and to evaluate the sensory properties of different blends of Aloe vera drink. The sensory qualities that were evaluated include color, taste, smell, appearance and feels. For each 200 ml of water, 14 g Aloe gel, 12.4 g Sugar, 3 Fresh mint leaves, 2 tbs of Fresh lemon juice and 0.37g of Ginger was used. The sensory qualities that were evaluated include color, taste, smell, appearance and feelings. The panelists were asked to record their observations and give marks to each parameter of different blends out of 10 on the sensory sheet. The most least liked Aloe blend was T4 that include Aloe gel, water, sugar and mint leaves. It was due to the strong smell and taste of mint leaves as mint leaves contain menthol. The overall acceptability of T8 was at the top as it contained complete blend of Aloe gel, mint leaves, fresh lemon juice and ginger. The potential of Aloe vera to be used in Food Industry and Pharmaceutical Industry has been increased. Aloe vera has many antibacterial, antifungal and antiviral active biological compounds which make it best for many disorders. An effort was made to make Aloe drink using the available resources and to develop awareness about its importance among the people.

Keywords: sensory qualities, pharmaceutical, immortality

Introduction:

The botanical name of Aloe vera is *Aloe barbadensis* Miller, it belongs to family Liliaceae. The name Aloe vera barbadensis has been specified for the true aloe, while the other aloes are normally known as Curacao aloe (Choi S and Chung M-H, 2003). Aloe vera has been used for its medicinal worth for several thousand years (J.L. Vogel zang, 2001). Its applications have been recorded in earliest cultures of several countries like Egypt, Greece, Rome and China. Its use is also mentioned in the Bible where it was taken as the plant of immortality (Saeed *et al*, 2004). Rajeswari, 2012 reported that Aloe is native to Africa and Mediterranean regions, but now it is spread to West Indies, India, China, Pakistan and other countries of Asia in 16th century (Ali and Qaiser, 2005). The name Aloe vera is normally considered that it comes from the Arabic word "Alloeh" meaning "Shining bitter substance" (Dandhof, 1987). There are about 350 different species of Aloe vera but generally all comes under the term "Aloe". The most commonly and abundantly used species of Aloe vera is *Aloe barbadensis* (Yagi *et al*, 2003).

Aloe vera is a perennial plant moreover it is an evergreen herb which is mainly concentrated in the tropical and sub-tropical regions of the world. It is a cactus like plant having large amount of pulp in its

leaves; it can easily survive in the hot and dry climates. The gel present in the leaves helps to store water and reduces the water loss and rate of evapo-transpiration (Sethi *et al*, 2012). Aloe is perennial which can reach up to the height of 30 cm. Leaves are fleshy, toothed at the margins and spirally arranged on the plant (Chaudhuri and Mukandan, 2001). The flower color of Aloe flower was described by Ali and Qaiser, 2005; they reported that in Pakistan and India its flowers are in reddish color while it has different color for the plants growing in Europe.

The leaves of Aloe vera consist of three important layers, a) the outer thick layer, b) a jelly like mucilage layer which is known as "aloes", c) the water storage area of the plant that is fillet. It has been revealed through research that the post harvest life of the Aloe vera leaves depends upon the methods of harvesting, handling and transportation. The Aloe vera leaf contains active biological compounds and their activity rate depends upon the handling and processing of Aloe vera leaves (Chandegara and Varshney, 2013).

Aloe vera gel has got much more attention as it is being used in the cosmetic industry for making creams, shampoos, lotions, in food industry for making different beverages and health drinks and in medical field for curing certain diseases (Hamman JH, 2008). It has been proved through previous study and

research that Aloe vera has more than 160 chemical compositions, from these 160 almost 70 chemical compositions are functional and active helps in maintaining the human health like anthraquinone compounds (Kong, 2003). Aloe vera contains 75 active constituents like enzymes, polysaccharides, essential and non-essential amino acids, beta-carotene, Vit C, E and B12, etc, and tannins together with their functional properties (Rashid *et al*, 2014). Similarly other compounds include polysaccharides, glucoside, lipids, organic acids, amino acids, enzymes, antibiotics and vitamins etc. fresh leaf of Aloe contain up to 96 % biologically neutral water of the total weight of leaf (Yan Li, 2009).

Now a day's Aloe gel extraction have got the research based and technical aspect as the gel is being used in different industries for variety of purposes. Chandegara and Varshney, 2013 discussed the different methods of gel extraction by hand filleting method. Mechanical filleting method is very common method to extract the aloe gel; this process takes place on conveyer belt equipped with blades (O'brien, 2005). In whole leaf processing method the tips and base is cut and then whole leaf is cut down in to pieces and treated with chemicals along with grinding to get the desirable constituents (Danhof, 2000). Roller squeezing method, Aloe vera leaf splitting method is also some important mechanical methods used for the extraction of Aloe vera leaf gel.

Aloe vera is now becoming an industrial crop as it is largely used in the food industry for the preparation of health drinks, beverages like tea, ice creams, dried juice, dessert and milk etc (Seoshin *et al*, 1995). Aloe vera juice is the most commonly used and the most of the Aloe gel is used to make Aloe drinks. All around the world different attempts have been done to make different recepies of Aloe drink which is good in taste, economical and also healthful. A blend of aloe vera, Anola and Ginger is being used to make Aloe drink in India (Sasi Kumar *et al*, 2013). Similarly Aloe drink is prepared by using different mixtures like Aloe vera gel, Ginger and Lemon juice extracts (Herlina, 2001). Now to increase the variety of tastes in the Aloe drinks different fruit juices are also used along with the Aloe gel to have the taste and medicinal value in one drink.

In the few recent years ethno botanical uses of plant material in the treatment of various diseases and ailments have got much attention. Chemical, biochemical, physio-chemical and biological studies on the Aloe vera plant have opened the new ways in medical, clinical and cosmetic fields. Aloe drink preparation with different mixtures that may include ginger, mint, lemon, stevia, anola or any other fruit is now become very common. As aloe drink have many medicinal uses and it is also used as energy drink so

different industries are trying to enhance their efficiency in making good and tasty juice with good shelf life.

The main objectives of this study are as under:

To develop the economical Aloe drink using available resources.

To evaluate the sensory properties of different blends of Aloe vera drink.

To evaluate the most liked and the best blend of Aloe drink.

Materials And Methods:

The research was conducted at Department of Horticulture, Pir Mehr Ali Shah Arid Agriculture University Rawalpind, Punjab, Pakistan in year 2013-14.

Harvesting Of Aloe Vera Leaves

Harvesting of aloe vera leaves is an important factor as the activity of biological compounds largely depends upon the method of harvesting. Aloe vera juice is used for treatment of several ailments along with the source of energy. So the biological activity of certain chemicals present in the leaf gel is very important. Micro-organisms infestation can reduce this activity if proper measurements are not adopted.

Under suitable environmental condition the healthy leaves with adequate amount of gel were harvested from the base of the plant using a sharp knife. The base of the leaf should remain intact and sealed while harvesting the leaf. Infestation of microorganisms increased if the leaves were harvested improperly. After harvesting of leaves, they were placed at suitable temperature and were processed within 4-5 hours in a room temperature. The leaves could be stored for about 3 weeks at if placed at appropriate temperature. Generally after every 7-8 weeks 3-4 healthy leaves of Aloe vera could be harvested.

Washing Of Harvested Aloe Leaves

Washing involved the removal of the dirt and other impurities from the leaf to get the pure gel with high level of active biological compounds. Harvested leaves were brought into the basin of the laboratory. Soft brushes were used to remove the adhered soil and dust particles from the harvested Aloe leaves. Then under the running tap water the leaves were properly washed. This process is repeated for several times until all the impurities were removed from the leaves. If the leaved were filled with the mud then it is suggested that they must be washed outside the laboratory before the final washing. The leaves were kept in a basket so that all the water drained away. The extraction of gel was done after the leaves were dried properly.

Extraction Of Leaf Gel

The gel extraction is the key step of Aloe leaves processing. The main constituent used for making Aloe drink is the Aloe gel, so the extraction method and leaf handling during the process is of supreme importance. Among various other gel extraction methods, the hand filleting method is commonly and frequently used. This method allows having the best quality and the maximum quantity if the Aloe vera leaves gel.

After washing and drying of the Aloe leaves, the gel extraction procedure begun with the peeling of the outer rind/peel. The tapering edges and the top and bottom portions of leaves were removed as it contains anthraquinone which would give bitter effect to the

Aloe drink. The peel was removed from the upper surface of leaf in such a way that minimum gel is attached to the leaves. The gel must be free from the green portion/rind of the leaves; extraction was done in a way to get the crystal clear gel. Once the upper layer of the rind was removed then gel was extracted by removing the lower layer of the rind. Another way to collect the gel was with the help of spoon. The extracted gel is collected and stored at proper temperature for later use. By hand filleting of the Aloe leaf, the leaf was used to get maximum quantity of gel and to make certain that the final product/blend contains all the nutrients that are present in the Aloe leaf.




Figure 1: Flow Chart Showing The Various Steps Involved In Making Aloe Drink.

Preparation Of Aloe Drink/Juice

Constituents Used

Aloe vera leaf gel, fresh leaves of mint, fresh lemon juice, ginger, sugar, water.

Types Of Blends:

- T1= Aloe gel + water
- T2= Aloe gel+ water + sugar
- T3= Aloe gel + water + sugar+ fresh lemon juice
- T4= Aloe gel+ water + sugar+ fresh mint leaves
- T5= Aloe gel+ water + sugar+ ginger
- T6= Aloe gel+ water + sugar+ fresh lemon juice + fresh mint leaves
- T7= Aloe gel+ water + sugar+ fresh lemon juice + ginger
- T8= Aloe gel+ water + sugar+ fresh lemon juice + fresh mint leaves + ginger

Quantities Of Ingredients

- Water = 200 ml
- Aloe gel = 14 g
- Sugar = 12.4 g
- Fresh mint leaves= 3 leaves
- Fresh lemon juice= 2 tbs
- Ginger = 0.37g

Preparation Of Aloe Juice

Freshly extracted gel of aloe, fresh lemon juice, fresh mint leaves and ginger were used to make Aloe drink. The lemon juice was obtained by removing seeds and squeezing the lemon, the mint leaves were chopped in to smaller pieces with knife and the ginger was first peeled and then chopped into smaller pieces. For all the above parameters the weight of Aloe gel, sugar and ginger, the lemon juice per liter and mint leaves were remain constant while the ingredients were different for various blends of Aloe. For different blends the ingredients were blended in a laboratory blender. After the preparation of each blend, the drink is poured in to the bottle of 1 liter and served to a group of individuals for the evaluation of sensory properties.

Sensory Quality Evaluation

In order to evaluate the sensory properties of Aloe drink a group of nine individuals were made including graduate students and some faculty members

of the department. The sensory qualities that were evaluated include color, taste, smell, appearance and feelings.

The panelists were asked to record their observations and give marks to each parameter of different blends out of 10 on the sensory sheet. The maximum likeness was shown by the number 10 and number 1 represents the minimum likeness.

Statistical Analysis

The experiment was one of the factorial setup in a entirely randomized design (CRD) with nine replications per treatment. Data were statistically analyzed by the analysis of variance (ANOVA) technique and the differences between different blends of Aloe vera were compared.

Results And Discussion

The colors of different blends were different ranging from transparent for T1 and T2 to greenish for T3 because of mint leaves and light yellow color for T4 as it contained lemon juice. The taste of different blends was also different according to various ingredients used. The marks given to the appearance was on the basis of overall look, transparency and reflection properties of the drink.

The comparative result of different parameters for specific blends was presented in (Figure 2). Color of T1 was best as it did not contain any constituent not even the sugar so transparency made it best in color. The color of T4 was least liked it was the blend containing mint leaves. Similarly the taste of T1 was least liked as it was without sugar so it did not have any taste, T6 had the best taste; it was the blend containing both lemon juice and mint leaves. Smell of T1 was not liked while T3 and T8 had equal level of likeness by smell. T1 was least liked by the appearance while the best blend by the appearance was T8. Feel is the overall level of satisfaction after having the blend; T4 had the minimum likeness in feel while T8 was at the top. T8 was the blend containing all the ingredients, the overall acceptability of T8 was at the top.

Table 1: Statistical Analysis Of The Data For Sensory Qualities Of Different Blends Of Aloe Vera.

Treatments	Color	Taste	Smell	Appearance	Feel
T1	7.1111 a	4.1111 b	4.4444 c	5.8889 a	5.0000 cd
T2	7.0000 a	7.1111 a	5.7778 abc	6.1111 a	6.6667 ab
T3	6.5556 a	7.7778 a	7.5556 a	5.6667 a	7.3333 ab
T4	2.7778 b	4.3333 b	5.7778 abc	6.2222 a	3.7778 d
T5	5.6667 a	6.0000 ab	5.4444 bc	6.5556 a	6.3333 bc
T6	6.3333 a	7.8889 a	6.8889 ab	6.5556 a	8.2222 a
T7	6.2222 a	7.0000 a	6.6667 ab	6.1111 a	7.0000 ab
T8	6.3333 a	7.6667 a	7.5556 a	7.4444 a	8.2222 a




Figure 2: Comparison Of Sensory Qualities For Each Aloe Blend.

Sensory qualities of different blends of Aloe vera beverage were determined and presented in (figure 3). It could be observed from the (figure 3) that appearance, color, taste, smell, appearance, feeling and overall acceptability of beverage varied among different blends. The most least liked Aloe blend was T4 that include Aloe gel, water, sugar and mint leaves. It may be due to the strong smell and taste of mint leaves. The overall acceptability of T8 was at the top as it contained complete blend of Aloe gel, mint leaves, fresh lemon juice and ginger. In this blend the

strong taste of mint leaves was overcome by the lemon juice and ginger was only used to give little taste so this was most liked by the panel. The blend without sugar was T1 was not much liked but it is best for the diabetic patients and used as a medicine. Mostly people don't like drinks without sugar and same is the case of T1 blend. As far as medicinal value of Aloe vera is concerned the blend T1 is most appropriate for the sugar patients. The strong taste of T4 was because of the presence of fresh mint leaves, mint leaves contain menthol up to 50% which gave strong taste to

Aloe drink. It was because of menthol that mint was used as mouth freshner and for soothing effects. (Alvi *et al*, 2001). The results of T6 showed that strong effect of mint leaves was reduced by the organic acids of lemon, mainly the citric acid. It also depends upon the quantity of lemon used but it has been observed

that citric acid of lemon lowers the effect of mint's menthol. Similarly the pungent taste of ginger is because of gingerols and shogaols which made the Aloe blend slightly pungent (Muhammad Majeed and Lakshami Prakash, 2007).

Table 2: Total Numbers (Out Of 90) For Each Parameter Of Different Blends Of Aloe Drink Given By A Group Of Nine Individuals.

Types Of Blends	Color	Taste	Smell	Appearance	Feeling
T1	64	32	40	53	45
T2	63	65	52	55	60
T3	59	70	68	51	65
T4	25	39	52	56	34
T5	51	54	49	59	57
T6	57	71	62	59	74
T7	56	63	60	55	63
T8	57	69	68	67	74




Figure 3: Sensory Quality Of Aloe Drink Blended With Aloe Vera, Lemon Juice, Mint Leaves And Ginger Juice.

Overall Acceptability

The overall acceptability of different Aloe blends was checked by adding the numbers given to different parameters of specific blend. For example, all the numbers for the parameters of blend T1 that was Color, Taste, Smell, Appearance and Feel were added to know that how much T1 was liked by the panel as compared to other blends. T4 got the least marks in over all acceptability it was the blend containing mint leaves. The second highest position got by the blend T6 it contained lemon juice and mint leaves to make a tasty Aloe blend. The blend that got the highest position in overall likeness was T8, as it contain the taste of min leaves, the citric acid of lemon juice made it slightly bitter and the minor taste of ginger made it overall a best blend.

Table 3: Total Numbers Given To Specific Parameters Of Different Blends.

Type of Blend	Overall Acceptability
T1	234
T2	295
T3	313
T4	206
T5	287
T6	323
T7	297
T8	335




Figure 4: Graphical View Of Overall Acceptability Of Different Aloe Blends.

Summary And Conclusion

The present study was conducted at Post Harvest Laboratory, Department of Horticulture, PMAS-Arid Agriculture University, Rawalpindi in year 2014 with the aim to make different blends of Aloe vera, due to the potential of Aloe vera to be used in Food Industry and Pharmaceutical Industry. Aloe vera has many antibacterial, antifungal and antiviral active biological compounds which make it best for many disorders.

Aloe vera has a long history as an herbal plant and is most popular for the treatment of various ailments. Aloe vera increases the availability of Vitamin C and E in the human body. It has also been proven that with the increase absorption of Vitamin C, Iron concentration in the body also been increased thus improving the health of people with iron deficiency. Similarly Aloe vera is also found helpful in wound healing, lowering blood sugar level and to cure

the cardiovascular disorders. Aloe vera is famous for proper digestion, blood circulation, for proper kidney functioning and maintaining the liver and colon function. It also has the ability to alkalize the digestive juices thus preventing the acidity of stomach. Researchers are going on to find out the effectiveness of Aloe vera in the curing and treatment of cancer and AIDS. Aloe vera is helpful in promoting the growth of non-cancerous cells.

In present investigation, efforts were made to develop different blends of Aloe drink and tried to evaluate the best one. Healthy and long leaves were suitable for the gel extraction as they contain large amount of leaf pulp. Care should be taken while extracting the gel, don't let the yellow substance that is anthraquinone to mix with the gel. Sensory qualities of all blends were carefully evaluated to find out the most favorite blend. An effort was made to make Aloe

drink using the available resources and to develop awareness about its importance among the people.

Recommendations

Different doses of Aloe vera gel can be used to evaluate its acceptability among different people. The storage life of different blends of Aloe drink can also be determined by using certain organic and chemical means.

Literature Cited

1. Ali, S. I. and M. Qaiser. 2005. Asphodelaceae, Flora of Pakistan. 211: 1-28. Department of Botany, University of Karachi and Missouri Botanical Press, Missouri Botanical Gardens, St. Louis, Missouri, USA.
2. Alvi, M.N., S. Ahmad and K. Rehman. 2001. Preparation of Menthol Crystals from Mint (*Mentha arvensis*). *Int. J. Agri. Biol.*, 3(4): 527–528.
3. Anonymous. 2008a. Post harvest processing of Aloe-vera leaves -a medicinal plant, All India Coordinated Research Project on Post harvest Technology (Indian Council of Agriculture Research), Annual Report, Bhubaneswar.
4. Boudreau, M.D. and F.A. Beland. 2006. An Evaluation of the Biological and Toxicological properties of Aloe barbadensis (miller), Aloe vera. *J. Environ. Sci. Health C Environ. Carcinog. Ecotoxicol. Rev.*, 24(1): 103-54.
5. Chandegara, V. K. and A. K. Varshney. 2014. Effect of Centrifuge Speed on Gel Extraction from Aloe Vera Leaves. *J. Food Process Technol.*, 5(1): 1000295.
6. Chandegara, V.K., and A.K. Varshney. 2013. Aloe vera L Processing and Products: A review. *Int. J. Med. Arom. Plant.*, 3(4), pp. 429-506, ISSN 2249-4340.
7. Chaudhuri, S. and U. Mukundan. 2001. Aloe vera M, Micropropagation and Characterization of its gel. *Phytomorphology*, 51:155-157.
8. Chauhan, O.P., P.S. Raju, F. Khanum and A.S. Bawa. 2007. Aloe vera- Therapeutic and food applications. *Indian Food Industry.*, 26(3): 43-51.
9. Choi, S. and M. H. Chung. 2003. A review on the relationship between Aloe vera components and their biological effects, *Seminars. In Integr. Medicine.*, 1:53–62.
10. Dagne, E., D. Bisrat, A. Viljoen and B.E. Van Wyk. 2000. Chemistry of Aloe species. *Current Organic Chemistry.*, 4(10), pp. 1055-1078(24).
11. Danhof, I.E. 2000. The Whole Leaf Advantage Excerpts <http://www.wholeleaf.com> Accessed.
12. Dandhof, I.E. 1987. Remarkable Aloe, Ommimedicus Press, Grand Prairie, Texas.
13. Grover, J.K., S. Yadav and V. Vats. 2002. Medicinal plants of India with anti-diabetic potential. *J. Ethnopharmacol.* 81(1): 81-100.
14. Hamman, J. H. 2008. Composition and applications of Aloe vera leaf gel., *Molecules* 13: 1599-1616.
15. Herlina, L. 2001. Incorporating private- public relationship into food technology development: creating added value to “Aloe vera” product SEAG symposium., Los Banos, Philippines.
16. Kumar, K. P. S., B. Debjit, Chiranjib and Biswajit. 2010. Aloe vera: A Potential Herb and its Medicinal Importance. *J. Chem. Pharm. Res.*, 2(1): 21-29.
17. Muhammad, Mujeeb and P. Lakshmi. 2007. Ginger (*Zingiber officinale*) Product Write Up. <http://www.sabinsa.com>.
18. Ni, Y., D. Turner, K.M. Yates and I. Tizard. 2004. Isolation and Characterization of Structural Components of Aloe vera L. leaf pulp. *Int. Immunophar.*, 4(14): 1745-55.
19. O'brien and Chantal. 2005. Physical and Chemical Characteristics of Aloe Gels. An unpublished M. sc. Thesis. University of Johannesburg.
20. Pankaj, K. S., D. G. Deen, S. Ritu, P. Priyanka, G. Sharmistha, K. S. Atul, K. Ajay and D. P. Kapil. 2013. Therapeutic and Medicinal Uses of Aloe Vera: a Review, *Pharmacology & Pharmacy.*, 4: 599-610.
21. Rajasekaran, S., K. Sivagnanam, K. Ravi and S. Subramanian. 2005. Antioxidant effect of Aloe vera gel extract in streptozotocin-induced diabetes in rats. *Pharmacol. Rep.* 57:90-96.
22. Rajeswari, R., M. Umadevi, C. S. Rahale, R. Pushpa, S. Selvavenkadesh, K. P. Sampath Kumar and D. Bhowmik. 2012. Aloe vera, The Miracle Plant Its Medicinal and Traditional Uses in India. *Journal of Pharmacognosy and Phytochemistry.*, 1(4): 118.
23. Ramachandra, C. T. and R. P. Srinivasa. 2008. Processing of Aloe Vera Leaf Gel: a Review. *American Journal of Agricultural and Biological Sciences.*, 3(2): 502-510.
24. Farhat, R., K. Tasneem, M. Q. Tahir, H. Sarfraz, N. Muhammad, A. Ammara and S. Z. Mahvish. 2014. Physico-chemical and sensory properties of orange marmalade supplemented with aloe vera powder. *J. Agric. Res.*, 2014, 52(4).
25. Saeed, M. A., A. Ishtiaq, Y. Uzma, A. Shazia, W. Amran, M. Saleem, and Nasir-ud-Din. 2004. Aloe Vera: A Plant of Vital Significance. *Quarterly Science Vision.*, Vol.9 No.1-2 (Jul - Dec, 2003) & 3-4 (Jan - Jun, 2004).
26. Sasi, K. R., R. C. Ramesh, P. K. Prodyut and C.P. Suresh. 2013. Development and Storage Studies of Therapeutic Ready to Serve (RTS)

- Made from Blend of Aloe vera, Aonla and Ginger Juice. *J. Food Process Technol.*, 4(6): 1000232.
27. Seoshin, Y., K. S. Lee, J. S. Lee and C. H. Lee. 1995. Preparation of Ghurt added with Aloe vera and its Quality Characteristics., *J. Korean Soc Food Nutr* 24:254-260.
 28. Sethi, J., A. Gupta, S. Sood, K. Dahiya, G. Singh and R. Gupta. 2012. Antioxidant Effect of Aloe vera in experimentally induced Diabetes Mellitus. *Int. J. Pharm. Sci.*, 3(8): 2522-2526.
 29. Stenhouse, J. 1851. On aloin, the Crystalline Cathartic principle of Barbadoes aloes. *Phil. Mag.* xxxvii: 481.
 30. Sushma, P., S. Alka and P. Devanand. 2013. Aloe vera: Composition, Processing and Medicinal Properties, *International journal of current discoveries and innovations.*, 2(2): 106-122.
 31. Vogelzang, J.L. 2001. "What You Need to Know About DietrySupplements", *Home Health Nurse*, 19, 50-52.
 32. Wang, Y.T. and K.J. Strong. 1995. Two-year study Monitoring several Physical and Chemical properties of field-grown Aloe barbadensis Miller leaves. *Subtropical Plant Sci.* 47:34-38.
 33. Xiangrong, K. and Z. Wang. 2003. Polysaccharide Chemical Structure and Biological activity of Aloe. *Herbal Medicine Abroad*.
 34. Yagi, A., A. Kabash, K. Mizuno, S. M. Moustafa, T. I. Khalifa and H. Tsuji. 2003. "Radical Scavenging Glycoprotein Inhibiting Cyclooxygenase-2 and Thromboxane A2 Syn-thase from Aloe vera Gel," *Planta Medica.*, 69(3), pp. 269-271.
 35. Yan, L. 2009. The Health Efficacy of Aloe and its Development and Utilization, *Asian Soil Science.*, 5(9).